

WGA Iso-lectin II WGA2, Isolectin D, WGA-B	
Molecular Weight	21,356Da [This is the MW of the unprocessed precursor]
Amino Acids Residue	213A.A. [This is the length of the unprocessed precursor]
Gene length	mRNA: 998bp ⁴⁾
Subunit	homodimer
Family	Chitin-binding type-1 domein
Accession No.(UniProtKB)	P02876
PDB ID (3D Structure)	2WGC (3'Sialyllactose) ³⁾ , 9WGA ³⁾
Metallic ion	None
Glycosylation Site	None
Cabos DB No.	N/A
WGA Isolectin III WGA3 (Fragment)	
Molecular Weight	18,756Da [This is the MW of the unprocessed precursor]
Amino Acids Residue	186A.A. [This is the length of the unprocessed precursor]
Gene length	mRNA: 898bp ⁶⁾
Subunit	homodimer
Family	Chitin-binding type-1 domein
Accession No.(UniProtKB)	P10969
PDB ID (3D Structure)	1K7T (GlcNAc β 1-6 Gal) ⁷⁾ , 1K7U (GlcNAc β 1-6GlcNAc) ⁷⁾ 1K7V (GlcNAc β 1-6Gal β 1-4Glc) ⁷⁾ , 1WGT ⁸⁾
Metallic ion	None
Glycosylation Site	None
Cabow DB No.	N/A

Reference

- 1) Nagata Y, Burger MM, *J. Biol. Chem.*,(1974), 249, 3116
- 2) Sugawara I, *RIKEN Rev.*,(1995), 8, 23
- 3) Wright CS, *J Mol Biol.*,(1990), 215, 635
- 4) Smith,J.J. , Raikhel,N.V. , *Plant Mol Biol.*,(1989), 13,601
- 5) Wright CS, Jaeger J,*J Mol Biol.*,(1993), 232, 620
- 6) Raikhel,N.V. , Wilkins,T.A., *Proc Natl Acad Sci U S A* 1987, 84 (19):6745-6749
- 7) Muraki M, Ishimura M, Harata K., *Biochim Biophys Acta.*,(2002), 1569,10
- 8) Harata K, Nagahora H, Jigami Y., *Acta Crystallogr D Biol Crystallogr.*,(1995), 51,1013